

References:

[Efficacy of an Andrographis paniculata composition for the relief of rheumatoid arthritis symptoms: a prospective randomized placebo-controlled trial.](#)

Burgos RA, Hancke JL, Bertoglio JC, Aguirre V, Arriagada S, Calvo M, Cáceres DD. Clin Rheumatol. 2009 Aug;28(8):931-46. doi: 10.1007/s10067-009-1180-5. Epub 2009 Apr 29.

[Andrographis paniculata decreases fatigue in patients with relapsing-remitting multiple sclerosis: a 12-month double-blind placebo-controlled pilot study.](#)

Bertoglio JC, Baumgartner M, Palma R, Ciampi E, Carcamo C, Cáceres DD, Acosta-Jamett G, Hancke JL, Burgos RA. BMC Neurol. 2016 May 23;16:77. doi: 10.1186/s12883-016-0595-2.

[\[Adalimumab plus methotrexate for the treatment of rheumatoid arthritis: a multi-center randomized, double-blind, placebo-controlled clinical study.\]](#)

Huang F, Zhang FC, Bao CD, Tao Y, Gu JR, Xu JH, Zhu P, Xu HJ, Zhang ZY, Zhao DB, Wu DH. Zhonghua Nei Ke Za Zhi. 2009 Nov;48(11):916-21. Chinese.

[Efficacy of tacrolimus in rheumatoid arthritis patients who have been treated unsuccessfully with methotrexate: a six-month, double-blind, randomized, dose-ranging study.](#)

Furst DE, Saag K, Fleischmann MR, Sherrer Y, Block JA, Schnitzer T, Rutstein J, Baldassare A, Kaine J, Calabrese L, Dietz F, Sack M, Senter RG, Wiesenhutter C, Schiff M, Stein CM, Satoi Y, Matsumoto A, Caldwell J, Harris RE, Moreland LW, Hurd E, Yocum D, Stampler DA. Arthritis Rheum. 2002 Aug;46(8):2020-8.

[Andrographis paniculata extract \(HMPL-004\) for active ulcerative colitis.](#)

Sandborn WJ, Targan SR, Byers VS, Ruddy DA, Mu H, Zhang X, Tang T. Am J Gastroenterol. 2013 Jan;108(1):90-8. doi: 10.1038/ajg.2012.340. Epub 2012 Oct 9.

[Use of visual analogue scale measurements \(VAS\) to assess the effectiveness of standardized Andrographis paniculata extract SHA-10 in reducing the symptoms of common cold. A randomized double blind-placebo study.](#)

Cáceres DD, Hancke JL, Burgos RA, Sandberg F, Wikman GK. Phytomedicine. 1999 Oct;6(4):217-23.

[Balneotherapy \(or spa therapy\) for rheumatoid arthritis.](#)

Verhagen AP, Bierma-Zeinstra SM, Boers M, Cardoso JR, Lambeck J, de Bie R, de Vet HC. Cochrane Database Syst Rev. 2015 Apr 11;(4):CD000518. doi: 10.1002/14651858.CD000518.pub2. Review.

[Double-blind, placebo-controlled, randomized, pilot clinical trial of ImmunoGuard--a standardized fixed combination of Andrographis paniculata Nees, with Eleutherococcus senticosus Maxim, Schizandra chinensis Bail. and Glycyrrhiza glabra L. extracts in patients with Familial Mediterranean Fever.](#)

Amaryan G, Astvatsatryan V, Gabrielyan E, Panossian A, Panosyan V, Wikman G. Phytomedicine. 2003 May;10(4):271-85.

[Radiographic, clinical, and functional outcomes of treatment with adalimumab \(a human anti-tumor necrosis factor monoclonal antibody\) in patients with active rheumatoid arthritis receiving concomitant methotrexate therapy: a randomized, placebo-controlled, 52-week trial.](#)

Keystone EC, Kavanaugh AF, Sharp JT, Tannenbaum H, Hua Y, Teoh LS, Fischkoff SA, Chartash EK.

Arthritis Rheum. 2004 May;50(5):1400-11.

[A comparative study of etanercept plus methotrexate and methotrexate alone in Taiwanese patients with active rheumatoid arthritis: a 12-week, double-blind, randomized, placebo-controlled study.](#)

Lan JL, Chou SJ, Chen DY, Chen YH, Hsieh TY, Young M Jr.
J Formos Med Assoc. 2004 Aug;103(8):618-23.

[Randomized double blind trial of an extract from the pentacyclic alkaloid-chemotype of uncaria tomentosa for the treatment of rheumatoid arthritis.](#)

Mur E, Hartig F, Eibl G, Schirmer M.
J Rheumatol. 2002 Apr;29(4):678-81.

[Golimumab in patients with active rheumatoid arthritis after treatment with tumour necrosis factor alpha inhibitors \(GO-AFTER study\): a multicentre, randomised, double-blind, placebo-controlled, phase III trial.](#)

Smolen JS, Kay J, Doyle MK, Landewé R, Matteson EL, Wollenhaupt J, Gaylis N, Murphy FT, Neal JS, Zhou Y, Visvanathan S, Hsia EC, Rahman MU; GO-AFTER study investigators..
Lancet. 2009 Jul 18;374(9685):210-21. doi: 10.1016/S0140-6736(09)60506-7. Epub 2009 Jun 26. Erratum in: Lancet. 2009 Oct 24;374(9699):1422.

[Golimumab, a human anti-tumor necrosis factor alpha monoclonal antibody, injected subcutaneously every four weeks in methotrexate-naive patients with active rheumatoid arthritis: twenty-four-week results of a phase III, multicenter, randomized, double-blind, placebo-controlled study of golimumab before methotrexate as first-line therapy for early-onset rheumatoid arthritis.](#)

Emery P, Fleischmann RM, Moreland LW, Hsia EC, Strusberg I, Durez P, Nash P, Amante EJ, Churchill M, Park W, Pons-Estel BA, Doyle MK, Visvanathan S, Xu W, Rahman MU.
Arthritis Rheum. 2009 Aug;60(8):2272-83. doi: 10.1002/art.24638. Erratum in: Arthritis Rheum. 2010 Oct;62(10):3005.

[Efficacy and tolerability of valdecoxib in treating the signs and symptoms of severe rheumatoid arthritis: a 12-week, multicenter, randomized, double-blind, placebo-controlled study.](#)

Gibofsky A, Rodrigues J, Fiechtner J, Berger M, Pan S.
Clin Ther. 2007 Jun;29(6):1071-85.

[A multicenter, double-blind, randomized, controlled phase III clinical trial of chicken type II collagen in rheumatoid arthritis.](#)

Wei W, Zhang LL, Xu JH, Xiao F, Bao CD, Ni LQ, Li XF, Wu YQ, Sun LY, Zhang RH, Sun BL, Xu SQ, Liu S, Zhang W, Shen J, Liu HX, Wang RC.
Arthritis Res Ther. 2009;11(6):R180. doi: 10.1186/ar2870. Epub 2009 Dec 1.

[Andrographis paniculata in the treatment of upper respiratory tract infections: a systematic review of safety and efficacy.](#)

Coon JT, Ernst E.
Planta Med. 2004 Apr;70(4):293-8. Review.

[A randomized double blind placebo controlled clinical evaluation of extract of Andrographis paniculata \(KalmCold\) in patients with uncomplicated upper respiratory tract infection.](#)

Saxena RC, Singh R, Kumar P, Yadav SC, Negi MP, Saxena VS, Joshua AJ, Vijayabalaji V, Goudar KS, Venkateshwarlu K, Amit A.
Phytomedicine. 2010 Mar;17(3-4):178-85. doi: 10.1016/j.phymed.2009.12.001. Epub 2010 Jan 25.

[A randomized, double-blind, multicenter, controlled clinical trial of cyclosporine plus chloroquine vs. cyclosporine plus placebo in early-onset rheumatoid arthritis.](#)

Miranda JM, Alvarez-Nemegyei J, Saavedra MA, Terán L, Galván-Villegas F, García-Figueroa J, Jara LJ, Barile L; GRECIA Group.
Arch Med Res. 2004 Jan-Feb;35(1):36-42.

[Randomized, double-blind, placebo-controlled, comparative study of human anti-TNF antibody adalimumab in combination with methotrexate and methotrexate alone in Taiwanese patients with active rheumatoid arthritis.](#)

Chen DY, Chou SJ, Hsieh TY, Chen YH, Chen HH, Hsieh CW, Lan JL.
J Formos Med Assoc. 2009 Apr;108(4):310-9. doi: 10.1016/S0929-6646(09)60071-1.

[A comparison of valdecoxib and naproxen in the treatment of rheumatoid arthritis symptoms.](#)

Williams GW, Kivitz AJ, Brown MT, Verburg KM.
Clin Ther. 2006 Feb;28(2):204-21

[Clinical improvement as reflected in measures of function and health-related quality of life following treatment with leflunomide compared with methotrexate in patients with rheumatoid arthritis: sensitivity and relative efficiency to detect a treatment effect in a twelve-month, placebo-controlled trial. Leflunomide Rheumatoid Arthritis Investigators Group.](#)

Tugwell P, Wells G, Strand V, Maetzel A, Bombardier C, Crawford B, Dorrier C, Thompson A.
Arthritis Rheum. 2000 Mar;43(3):506-14. Erratum in: Arthritis Rheum 2000 Jun;43(6):1345.

[Combination therapy with cyclosporine and methotrexate in severe rheumatoid arthritis. The Methotrexate-Cyclosporine Combination Study Group.](#)

Tugwell P, Pincus T, Yocum D, Stein M, Gluck O, Kraag G, McKendry R, Tesser J, Baker P, Wells G.
N Engl J Med. 1995 Jul 20;333(3):137-41.

[Efficacy and safety of willow bark extract in the treatment of osteoarthritis and rheumatoid arthritis: results of 2 randomized double-blind controlled trials.](#)

Biegert C, Wagner I, Lüdtke R, Kötter I, Lohmüller C, Günaydin I, Taxis K, Heide L.
J Rheumatol. 2004 Nov;31(11):2121-30.

[Dose escalation of parenteral methotrexate in active rheumatoid arthritis that has been unresponsive to conventional doses of methotrexate: a randomized, controlled trial.](#)

Lambert CM, Sandhu S, Lochhead A, Hurst NP, McRorie E, Dhillon V.
Arthritis Rheum. 2004 Feb;50(2):364-71.

[Tramadol 37.5-mg/acetaminophen 325-mg combination tablets added to regular therapy for rheumatoid arthritis pain: a 1-week, randomized, double-blind, placebo-controlled trial.](#)

Lee EY, Lee EB, Park BJ, Lee CK, Yoo B, Lim MK, Shim SC, Sheen DH, Seo YI, Kim HA, Baek HJ, Song YW.
Clin Ther. 2006 Dec;28(12):2052-60.

[Randomized double blind trial of an ayurvedic plant derived formulation for treatment of rheumatoid arthritis.](#)

Chopra A, Lavin P, Patwardhan B, Chitre D.
J Rheumatol. 2000 Jun;27(6):1365-72.

[Effects of abatacept in patients with methotrexate-resistant active rheumatoid arthritis: a randomized trial.](#)

Kremer JM, Genant HK, Moreland LW, Russell AS, Emery P, Abud-Mendoza C, Szechinski J, Li T, Ge Z, Becker JC, Westhovens R.
Ann Intern Med. 2006 Jun 20;144(12):865-76.

[Andrographis paniculata in the symptomatic treatment of uncomplicated upper respiratory tract infection: systematic review of randomized controlled trials.](#)

Poolsup N, Suthisisang C, Prathanturarug S, Asawamekin A, Chanchareon U.
J Clin Pharm Ther. 2004 Feb;29(1):37-45. Review.

[Evaluation of the safety of rituximab in combination with a tumor necrosis factor inhibitor and methotrexate in patients with active rheumatoid arthritis: results from a randomized controlled trial.](#)

Greenwald MW, Shergy WJ, Kaine JL, Sweetser MT, Gilder K, Linnik MD.
Arthritis Rheum. 2011 Mar;63(3):622-32. doi: 10.1002/art.30194.

[A randomised controlled trial of etanercept and methotrexate to induce remission in early inflammatory arthritis: the EMPIRE trial.](#)

Nam JL, Villeneuve E, Hensor EM, Wakefield RJ, Conaghan PG, Green MJ, Gough A, Quinn M, Reece R, Cox SR, Buch MH, van der Heijde DM, Emery P.
Ann Rheum Dis. 2014 Jun;73(6):1027-36. doi: 10.1136/annrheumdis-2013-204882. Epub 2014 Mar 11.

[Methotrexate for treating rheumatoid arthritis.](#)

Lopez-Olivo MA, Siddhanamatha HR, Shea B, Tugwell P, Wells GA, Suarez-Almazor ME.
Cochrane Database Syst Rev. 2014 Jun 10;(6):CD000957. doi: 10.1002/14651858.CD000957.pub2. Review.

[A randomized, double-blinded, placebo-controlled study of the effect of a combination of lemon verbena extract and fish oil omega-3 fatty acid on joint management.](#)

Caturla N, Funes L, Pérez-Fons L, Micol V.
J Altern Complement Med. 2011 Nov;17(11):1051-63. doi: 10.1089/acm.2010.0410.

[Combination of Mangifera indica L. extract supplementation plus methotrexate in rheumatoid arthritis patients: a pilot study.](#)

López Mantecón AM, Garrido G, Delgado-Hernández R, Garrido-Suárez BB.
Phytother Res. 2014 Aug;28(8):1163-72. doi: 10.1002/ptr.5108. Epub 2013 Dec 17.

[Everolimus in patients with rheumatoid arthritis receiving concomitant methotrexate: a 3-month, double-blind, randomised, placebo-controlled, parallel-group, proof-of-concept study.](#)

Bruyn GA, Tate G, Caeiro F, Maldonado-Cocco J, Westhovens R, Tannenbaum H, Bell M, Forre O, Bjerneboe O, Tak PP, Abeywickrama KH, Bernhardt P, van Riel PL; RADD Study Group.
Ann Rheum Dis. 2008 Aug;67(8):1090-5. Epub 2007 Nov 23.

[Assessment of comparative pain relief and tolerability of SKI306X compared with celecoxib in patients with rheumatoid arthritis: a 6-week, multicenter, randomized, double-blind, double-dummy, phase III, noninferiority clinical trial.](#)

Song YW, Lee EY, Koh EM, Cha HS, Yoo B, Lee CK, Baek HJ, Kim HA, Suh YI, Kang SW, Lee YJ, Jung HG.
Clin Ther. 2007 May;29(5):862-73.

[Comparison of Tripterygium wilfordii Hook F with methotrexate in the treatment of active rheumatoid arthritis \(TRIFRA\): a randomised, controlled clinical trial.](#)

Lv QW, Zhang W, Shi Q, Zheng WJ, Li X, Chen H, Wu QJ, Jiang WL, Li HB, Gong L, Wei W, Liu H, Liu AJ, Jin HT, Wang JX, Liu XM, Li ZB, Liu B, Shen M, Wang Q, Wu XN, Liang D, Yin YF, Fei YY, Su JM, Zhao LD, Jiang Y, Li J, Tang FL, Zhang FC, Lipsky PE, Zhang X. *Ann Rheum Dis.* 2015 Jun;74(6):1078-86. doi: 10.1136/annrheumdis-2013-204807. Epub 2014 Apr 14.

[A randomized trial of hydroxychloroquine in early rheumatoid arthritis: the HERA Study.](#)

[No authors listed]

Am J Med. 1995 Feb;98(2):156-68.

[Concomitant leflunomide therapy in patients with active rheumatoid arthritis despite stable doses of methotrexate. A randomized, double-blind, placebo-controlled trial.](#)

Kremer JM, Genovese MC, Cannon GW, Caldwell JR, Cush JJ, Furst DE, Luggen ME, Keystone E, Weisman MH, Bensen WM, Kaine JL, Ruderman EM, Coleman P, Curtis DL, Kopp EJ, Kantor SM, Waltuck J, Lindsley HB, Markenson JA, Strand V, Crawford B, Fernando I, Simpson K, Bathon JM.

Ann Intern Med. 2002 Nov 5;137(9):726-33.

[omega-3 Fatty acids infusions as adjuvant therapy in rheumatoid arthritis.](#)

Bahadori B, Uitz E, Thonhofer R, Trummer M, Pestemer-Lach I, McCarty M, Krejs GJ.

JPEN J Parenter Enteral Nutr. 2010 Mar-Apr;34(2):151-5. doi: 10.1177/0148607109342130.

[Golimumab, a new human anti-tumor necrosis factor alpha antibody, administered intravenously in patients with active rheumatoid arthritis: Forty-eight-week efficacy and safety results of a phase III randomized, double-blind, placebo-controlled study.](#)

Kremer J, Ritchlin C, Mendelsohn A, Baker D, Kim L, Xu Z, Han J, Taylor P.

Arthritis Rheum. 2010 Apr;62(4):917-28. doi: 10.1002/art.27348. Erratum in: *Arthritis Rheum.* 2010 Oct;62(10):3130.

Nakamura H, Nishioka K. Effects of glucosamine/chondroitin supplement on osteoarthritis: involvement of PGE2 and YKL-40. *Jpn J Joint Dis.* 2002;11:175-84.

Asari A, Kanemitsu T, Kurihara H. Oral administration of high molecular weight hyaluronan (900 kDa) controls immune system via Toll-like receptor 4 in the intestinal epithelium. *J Biol Chem.*

2010;285:24751-8. doi: 10.1074/jbc.M110.104950

Hatayama T, Nagano M, Yamaguchi N, Kumagai S, Ohnuki K. The effect of a supplement on knee pain and discomfort evaluated by visual analogue scale(VAS): a randomized, double-blind, placebo-controlled study. *Kenko-shien.* 2008;10:13-7.

Jensen GS, Attridge VL, Lenninger MR, Benson KF. Oral intake of a liquid high-molecular-weight hyaluronan associated with relief of chronic pain and reduced use of pain medication: results of a randomized, placebo-controlled double-blind pilot study. *J Med Food.* 2015;18:95-101. doi: 10.1089/jmf.2013.0174.